

February 2, 2008

Heat can flow from cold to hot in Microcanonical Thermodynamics of finite systems and the microscopic origin of phase transitions

D.H.E. Gross

*Hahn-Meitner Institute and Freie Universität Berlin,
Fachbereich Physik.
Glienickestr. 100
14109 Berlin, Germany**

Microcanonical Thermodynamics [1] allows the application of Statistical Mechanics to finite and even small systems. As surface effects cannot be scaled away, one has to be careful with the standard arguments of splitting a system into two or bringing two systems into thermal contact with energy or particle exchange: Not only the volume part of the entropy must be considered. The addition of any other macroscopic constraint like a dividing surface, or the enforcement of gradients of the energy/particle reduce the entropy. As will be shown here, when removing such constraint in regions of a negative heat capacity, the system may even relax under a flow of heat *against the temperature slope*. Thus Clausius formulation of the Second Law: "Heat always flows from hot to cold" can be violated. However, the Second Law is still satisfied and the total Boltzmann-entropy is rising. In the final chapter the general microscopic mechanism leading to the convexity of the microcanonical entropy at phase separation is discussed. This is explained for the liquid-gas and the solid-liquid transition.

I. INTRODUCTION

In conventional (extensive) thermodynamics the thermal equilibrium of two systems is established by bringing them into thermal contact which allows free energy exchange[12]. Equilibrium is established when the total entropy

$$S_{total} = S_1 + S_2 \quad (1)$$

is maximal:

$$dS_{total} = dS_1 + dS_2 = 0. \quad (2)$$

Under an energy flux $\Delta E_{2 \rightarrow 1}$ from $2 \rightarrow 1$ the total entropy changes by

$$\Delta S_{total} = (\beta_1 - \beta_2) \Delta E_{2 \rightarrow 1} \quad (3)$$

$$\beta = dS/dE = \frac{1}{T} \quad (4)$$

Consequently, a maximum of $S_{total}(E = E_1 + E_2)$ will be approached when

$$\text{sign}(\Delta S_{total}) = \text{sign}(T_2 - T_1) \text{sign}(\Delta E_{2 \rightarrow 1}) > 0 \quad (5)$$

From here Clausius' first formulation of the Second Law follows: "Heat always flows from hot to cold". Essential for this conclusion is the *additivity* of S under the split (eq.1). Temperature is an appropriate control parameter for extensive systems.

II. STABILITY AGAINST SPONTANEOUS ENERGY-GRADIENTS

For a small or a very large self-gravitating system, c.f. [1], additivity and extensivity of S , and also of E , is not given. In fact this is the main reason to develop this new and extended version of thermodynamics. At phase separation the microcanonical caloric curve $T(E)$ is *backbending*. Here the heat capacity $C = -(\frac{\partial S}{\partial E})^2 / \frac{\partial^2 S}{\partial E^2}$ becomes *negative*, the curvature $\frac{\partial^2 S}{\partial E^2}$ is positive, and $S(E)$ is *convex*. This is the *generic signal of a phase transition of first order* [1]. The Potts-model illuminates in a most simple example the occurrence of a backbending caloric curve [2][13]. A typical plot of $s(e) = S(E = Ne)/N$ in the region of phase separation is shown in fig(1). Section III discusses the general microscopic reasons for the convexity.

Let us split the system into two pieces $a+b$ by a dividing surface, with half the number of particles each. The dividing surface is purely geometrical. It exists only as long as the two pieces can be distinguished by their different energy/particle e_a and e_b . If the constraint on the difference $e_b - e_a$ is fully relaxed and $e_b - e_a$ can fluctuate freely at fixed $e_2 = (e_a + e_b)/2$, the dividing surface is assumed to have no further physical effect on the system. Constraining the energy-difference $e_b - e_a = \Delta e$ between the two, reduces the number of free, unconstrained degrees of freedom and *reduces* the entropy by $-\Delta \Sigma$.

In the convex region (upwards concave like $y = x^2$) of $s(e)$ this becomes especially evident: at $e = e_2 \sim e_{min}$, $e_a = e_2 - \Delta e/2$ and $e_b = e_2 + \Delta e/2$, the simple algebraic sum of the individual entropies $s_{sum} = (s_a + s_b)/2$ leads to $s_{sum} \geq s_2$. However, e^{S_2} is already the sum over

*Electronic address: gross@hmi.de;
URL: <http://www.hmi.de/people/gross/>




FIG. 1: Potts model ($q = 10$) on a 50×50 -lattice with periodic boundary conditions in the region of phase separation. At the energy e_1 per lattice point the system is in the pure ordered phase, at e_3 in the pure disordered phase. At e_a little above e_1 the temperature $T_a = 1/\beta$ is higher than T_2 and even more than T_b at e_b a little below e_3 . At e_a the system separates into bubbles of disordered phase embedded in the ordered phase or at e_b into droplets of ordered phase within the disordered one. If we combine two equal systems: one with the energy per lattice site $e_a = e_1 + \Delta e$ and at the temperature T_a , the other with the energy $e_b = e_3 - \Delta e$ and at the temperature $T_b < T_a$, and allowing for free energy exchange, then the systems will equilibrate at energy e_2 with a rise of its entropy. The temperature T_a drops (cooling) and energy (heat) flows (on average) from $b \rightarrow a$. I.e.: Heat flows from cold to hot! Clausius formulation of the Second Law is violated. This is well known for self-gravitating systems. However, this is not a peculiarity of only gravitating systems! It is the generic situation at phase separations within classical thermodynamics even for systems with short-range coupling and *has nothing to do with long range interactions*.

all possible configurations at e_2 including the split ones with $e_b - e_a = \Delta e$. Evidently, the enforced energetic split of the system into two pieces with prescribed specific energies $e_b > e_a$ destroys configurations with different or vanishing $e_b - e_a$. This reduces the partition sum. It costs some additional "surface" or constraining entropy $\Delta\Sigma > 0$ to maintain the imbalance of the energy, such that $S_2 \geq S_{sum} - \Delta\Sigma$.

From this little exercise we learn that the microcanonical entropy per particle $s(e, N, \sigma)$ of a finite system depends not only on the energy per particle $e = E/N$ and possibly still on the particle number N (for a non-extensive system) but also on the boundary (the container), or any other non-extensive external constraint, via $\Sigma = N\sigma$. If one splits a given configuration like in eq.(1) one must not forget the change $-\Delta\Sigma$ in the surface entropy.

The correct entropy balance, before and after establishing the energetic split $e_b > e_a$ of the system, is

$$s_{after} - s_{before} = \frac{s_a + s_b}{2} - s_2 - \Delta\sigma \leq 0 \quad (6)$$

even though the sum of the first two terms is positive.

In the inverse direction: Only by *relaxing* the constraint and by allowing, on average, for an energy-flux ($\Delta e_{b \rightarrow a} > 0$) *opposite to* $T_a - T_b > 0$, *against the temperature-gradient (slope)*, but in the direction of the energy-slope, an increase of $s_{total} \rightarrow s_2$ is obtained. Then, on average both energies e_a and e_b become equal $= e_2$ and $\Delta\sigma = 0$. Or in other words, starting with an energy difference $\Delta e = e_b - e_a \geq 0$ the allowance of unconstrained energy fluctuations around $\Delta e = 0$ between the two subsystems implies the disappearance of $\Delta\sigma$ and in general leads to an increase of the total entropy. This is consistent with the naive picture of an energy *equilibration*. Thus Clausius' "Energy flows always from hot to cold" is violated. Of course this shows again that *unlike to extensive thermodynamics the temperature is not an appropriate control parameter in non-extensive systems*.

In the thermodynamic limit $N \rightarrow \infty$ of a system with short-range coupling $\Delta\Sigma \sim N^{2/3}$, $\Delta\Sigma/N = \Delta\sigma \propto N^{-1/3}$ must go to 0 due to van Hove's theorem.

III. THE ORIGIN OF THE CONVEXITY OF $S(E)$ AT PHASE SEPARATION

Here I discuss the general microscopic mechanism leading to the appearance of a convex intruder in $S(E, \mathcal{V}, \dots)$. This is the signal of phase transitions of first order and of phase-separation within the microcanonical ensemble.

I assume the system is classical and obeys the Hamiltonian:

$$H = \sum_{i \leq j=1, N} \left\{ \frac{p_i^2}{2m} \delta_{i,j} + V^{int}(\vec{r}_i - \vec{r}_j) \right\} \quad (7)$$

In this case the system is controlled by energy and volume.

A. Liquid-gas transition

The microcanonical sum of states or partition sum is:

$$\begin{aligned}
W(E, N, \mathcal{V}) &= \frac{1}{N!(2\pi\hbar)^{3N}} \int_{\mathcal{V}^N} d^{3N} \vec{r}_i \int d^{3N} \vec{p}_i \delta[E - \sum_i^N \frac{\vec{p}_i^2}{2m_i} - V^{int}\{\vec{r}_i\}] \\
&= \frac{\mathcal{V}^N (E - E_0)^{(3N-2)/2} \prod_1^N m_i^{3/2}}{N! \Gamma(3N/2) (2\pi\hbar^2)^{3N/2}} \int_{\mathcal{V}^N} \frac{d^{3N} r_i}{\mathcal{V}^N} \left(\frac{E - V^{int}\{\vec{r}_i\}}{E - E_0} \right)^{(3N-2)/2} \quad (8)
\end{aligned}$$

$$\begin{aligned}
&= W^{id-gas}(E - E_0, N, \mathcal{V}) \times W^{int}(E - E_0, N, \mathcal{V}) \\
&= e^{[S^{id-gas} + S^{int}]} \quad (9)
\end{aligned}$$

$$W^{id-gas}(E, N, \mathcal{V}) = \frac{\mathcal{V}^N E^{(3N-2)/2} \prod_1^N m_i^{3/2}}{N! \Gamma(3N/2) (2\pi\hbar^2)^{3N/2}} \quad (10)$$

$$W^{int}(E - E_0, N, \mathcal{V}) = \int_{\mathcal{V}^N} \frac{d^{3N} r_i}{\mathcal{V}^N} \left(1 - \frac{V^{int}\{\vec{r}_i\} - E_0}{E - E_0} \right)^{(3N-2)/2} \quad (11)$$

$\mathcal{V}$ is the spatial volume. $E_0 = \min V^{int}\{\vec{r}\}$ is the energy of the ground-state of the system.

The separation of $W(E, N, \mathcal{V})$ into W^{id-gas} and W^{int} is the microcanonical analogue of the split of the canonical partition sum into a kinetic part and a configuration part:

$$Z(T) = \frac{\mathcal{V}^N}{N!} \left(\frac{mT}{2\pi\hbar^2} \right)^{3N/2} \int \frac{d^{3N} r_i}{\mathcal{V}^N} e^{-\frac{V^{int}\{\vec{r}_i\}}{T}} \quad (12)$$

In the thermodynamic limit the order parameter of the (homogeneous) liquid-gas transition is the density. The transition is linked to a condensation of the system towards a larger density controlled by pressure. For a finite system we expect the analogue. However, here controlled by the constant available system volume $\mathcal{V}$. If it is larger than the eigen-volume $\mathcal{V}_0$ of the condensate, the system does not fill the volume $\mathcal{V}$ at low E . $N - 1$ internal coordinates are limited to $\mathcal{V}_0$. Only the center of mass of the droplet can move freely in $\mathcal{V}$. The system does not fill the 3N-configuration space $\mathcal{V}^N$. Only a stripe of the width $\mathcal{V}_0^{1/3}$ in $3(N - 1)$ dimensions of the total 3-N dim space and with the length $\mathcal{V}^{1/3}$ in the remaining 3 dimensions of center of mass motion is populated. *The system is non-homogeneous even though it is equilibrated and, at low energies, internally in the single liquid phase.* It is not characterized by an intensive homogeneous density. In fact, $W^{int}(E - E_0, N, \mathcal{V})$ can be written as:

$$W^{int}(E - E_0, N, \mathcal{V}) = \left[\frac{\mathcal{V}(E)}{\mathcal{V}} \right]^N \leq 1 \quad (13)$$

$$[\mathcal{V}(E)]^N \stackrel{\text{def}}{=} \int_{\mathcal{V}^N} d^{3N} r_i \left(1 - \frac{V^{int}\{\vec{r}_i\} - E_0}{E - E_0} \right)^{(3N-2)/2} \quad (14)$$

$$S^{int}(E - E_0, N, \mathcal{V}) = N \ln \left[\frac{\mathcal{V}(E)}{\mathcal{V}} \right] \leq 0 \quad (15)$$

In eqs.(11,14) the integration is only over regions in 3N-dim coordinate space where $V^{int}\{\vec{r}_i\}$ is smaller than E , where the big brackets are positive. The volume $\mathcal{V}^N(E) \leq \mathcal{V}^N$ is the accessible part of the spatial volume outside of the forbidden 3N-dim regions with $V^{int}\{r_i\} > E$. I.e. the eigen-volume of the condensate (droplets) at the given energy. $\mathcal{V}^N(E)$ has the limiting values:

$$[\mathcal{V}(E)]^N := \begin{cases} \mathcal{V}^N & \text{for } E \text{ in the gas phase} \\ \mathcal{V}_0^{N-1} \mathcal{V} & \text{for } E = E_0 \end{cases}$$

So that $W^{int}(E - E_0, N, \mathcal{V})$ and $S^{int}(E - E_0, N, \mathcal{V})$ have the property:

$$\begin{aligned}
W^{int}(E - E_0) &\leq 1, \Rightarrow S^{int}(E - E_0, N) \leq 0 \\
&\rightarrow \begin{cases} 1 & E \gg V^{int} \\ \left[\frac{\mathcal{V}_0}{\mathcal{V}} \right]^{(N-1)} & E \rightarrow E_0 \end{cases} \quad (16)
\end{aligned}$$

$$S^{int}(E - E_0) \rightarrow \begin{cases} 0 & E \gg V^{int} \\ \ln \left\{ \left[\frac{\mathcal{V}_0}{\mathcal{V}} \right]^{N-1} \right\} < 0 & E \rightarrow E_0 \end{cases} \quad (17)$$

All physical details are encrypted in $W^{int}(E - E_0, N, \mathcal{V})$ or $S^{int}(E - E_0, N, \mathcal{V})$ alias $N \ln[\mathcal{V}(E)]$, c.f. eqs.(13–17): If the energy is high the detailed structure of $V^{int}\{\vec{r}_i\}$ is unimportant $W^{int} \approx 1$, $S^{int} \approx 0$. The system behaves like an ideal gas and fills the volume $\mathcal{V}$. At sufficiently low energies only the minimum of $V^{int}\{\vec{r}_i\}$ is explored by $W^{int}(E - E_0, N, \mathcal{V})$. The system is in a condensed, liquid drop (perhaps several) moving freely inside the empty larger volume $\mathcal{V}$.

One can guess the general form of $N \ln[\mathcal{V}(E)]$: Near the groundstate $E \gtrsim E_0$ it must be flat $\approx (N - 1) \ln[\mathcal{V}_0] + \ln[\mathcal{V}]$ because the liquid drop has some eigen-volume $\mathcal{V}_0$ in which *each* particle can move. With rising energy $\ln[\mathcal{V}(E)]$ rises up to the point (E_{trans}) where it is possible that the drop fissions into two and $N \ln[\mathcal{V}(E)] \approx$

$(N-2)\ln[\mathcal{V}_0]+2\ln[\mathcal{V}]$. I.e. here is a first "jump" upwards of $S^{int}(E)$ by $\ln\{\frac{\mathcal{V}}{\mathcal{V}_0}\}$. Later further such "jumps" may follow. Each of these "jumps" induce a convex upwards bending also of the total entropy $S(E)$ (eq.9). Each is connected to a bifurcation and bimodality of $e^{S(E)-E/T}$ and the phenomenon of *phase-separation*. Of course in the conventional canonical picture for a large number of particles this is washed out and leads to the familiar Yang-Lee singularity of the liquid to gas phase transition. In the microcanonical ensemble this is analogue to the phenomenon of *multifragmentation* in nuclear systems [1, 3]. This, in contrast to the mathematical Y-L theorem, *physical* microscopic explanation of the liquid to gas phase transition shows the advantage of the microcanonical statistics on the $3N$ -coordinate space as presented here.

B. Solid-liquid transition

In contrast to the liquid phase, in the crystal phase a molecule can only move locally within its lattice cell of the size d^3 instead of the whole volume $\mathcal{V}_0$ of the condensate. I.e. in eq.(17) instead we have $S^{int} \rightarrow \ln\{[\frac{d^3}{\mathcal{V}_0}]^{N-1}\}$.

C. Summary of section III

The gas-liquid transition is linked to the transition from uniform filling of the container volume $\mathcal{V}$ by the

gas to the *smaller eigen-volume* of the system $\mathcal{V}_0$ in its condensed phase where the system is *non-homogeneous* (some liquid drops inside the larger *empty* volume $\mathcal{V}$). First $N-1$, later at higher energies less and less degrees of freedom condensate into the drop. First one, then more and more dofs. explore the larger container volume $\mathcal{V}$ leading to upwards jumps (convexities) of $S^{int}(E)$. The volume of the container controls how close one is to the critical end-point of the transition, where phase-separation disappears. Towards the critical end-point, i.e. with smaller $\mathcal{V}$ the jumps $\ln[\mathcal{V}] - \ln[\mathcal{V}_0]$ become smaller and smaller. In the case of the solid-liquid transition the external volume $\mathcal{V}$ of the container has no influence on $S^{int}(E)$. The entropy jumps are by $\Delta S^{int} \propto \ln \mathcal{V}_0 - \ln d^3$.

At the surface of the drops $V^{int} > E_0 = \min V^{int}$, i.e. the surface gives a *negative* contribution to S^{int} in eq.(14) and to S at energies $E \gtrsim E_0$, as was similarly assumed in section (II) and explicitly in eq.(6).

IV. ACKNOWLEDGEMENT

I am grateful to J. Möller for insistent, therefore helpful, discussions.

-
- [1] D.H.E. Gross. *Microcanonical thermodynamics: Phase transitions in "Small" systems*, volume 66 of *Lecture Notes in Physics*. World Scientific, Singapore, 2001.
 - [2] D.H.E. Gross, A. Ecker, and X.Z. Zhang. Microcanonical thermodynamics of first order phase transitions studied in the potts model. *Ann. Physik*, 5:446–452, 1996, and <http://xxx.lanl.gov/abs/cond-mat/9607150>.
 - [3] D.H.E. Gross. Microcanonical thermodynamics and statistical fragmentation of dissipative systems — the topological structure of the n-body phase space. *Physics Reports*, 279:119–202, 1997.
 - [4] L. G. Moretto, L. Phairand J. B. Elliot and, and G. J. Wozniak. Negative heat capacities and first order phase transitions in nuclei. *Phys. Rev.C*, 66:041601(R),nucl-theor/0208024, 2002.
 - [5] D.H.E. Gross. Statistical decay of very hot nuclei, the production of large clusters. *Rep.Progr.Phys.*, 53:605–658, 1990.
 - [6] M. D'Agostino, F. Gulminelli, Ph. Chomaz, M. Bruno, F. Cannata, R. Bougault, F. Gramegna, I. Iori, N. le Neindre, G.V. Margagliotti, A. Moroni, and G. Vannini. Negative heat capacity in the critical region of nuclear fragmentation: an experimental evidence of the liquid-gas phase transition. *Phys.Lett.B*, 473:219–225, 2000.
 - [7] B. Borderie. Dynamics and thermodynamics of the liquid-gas phase transition in hot nuclei studied with the Indra array. *J.Phys.G: Nucl.Part.Phys.*, 28:R212–R227, 2002.
 - [8] M. Schmidt, R. Kusche, T. Hippler, J. Donges, W. Kornmüller, B. von Issendorff, and H. Haberland. Negative heat capacity for a cluster of 147 sodium atoms. *Phys.Rev.Lett.*, 86:1191–1194, 2001.
 - [9] M. Schmidt and H. Haberland. Caloric curve across the liquid-to-gas change for sodium clusters. *Phys.Rev.Lett.*, 87:203402–1–4, 2001.
 - [10] D.H.E. Gross. Comment on "Negative heat and first order phase transitions in nuclei" by Moretto et al. <http://arXiv.org/abs/nucl-th/0208034>, 2002).
 - [11] F. Gulminelli and Ph. Chomaz. Comment on "Negative heat capacities and first order phase transitions in nuclei" by L.G. Moretto et al. *Caen preprint, LPCC 99-17*, nucl-th/0304058, 2003.
 - [12] The first two sections discuss mainly systems that have no other macroscopic (extensive) control parameter besides the energy. E.g. the particle density is not changed and there are no chemical reactions.
 - [13] This backbending of $T(e)$ has nothing to do with the periodic boundary condition used here, as claimed erroneously by Moretto et.al [4]. Many realistic microcanonical calculations for nuclear fragmentation as also for atomic-cluster fragmentation, c.f. [1, 5], show the back-

bending *without using periodic boundaries*. These are verified by numerous experimental data in nuclear physics c.f. [6, 7] and cluster physics [8, 9]. The errors of the

above paper by Moretto et. al. are commented in more detail also in [10, 11]